



7. Weissman, M.M., Verdelli, H., Gamerooff, M.J., Bledsoe, S.E., Betts K, et al. (2006). National Survey of Psychotherapy training in psychiatry, psychology, and social work. *Archives of General Psychiatry*, 63, 925–934.

Partnering Hospital Services and Community Therapists for People with Borderline Personality Disorder: Dialectical Behaviour Therapy 'Lite'

Sarah Birnie, Ph.D., C.Psych.

Outpatient Mental Health, The Ottawa Hospital, Civic Campus

References

- Cohen, K. (2012, February). *Needs, demands and supply in professional psychology or how well is psychology prepared to do what with whom?* Colloquia presented at Psychology Day, The Ottawa Hospital, Ottawa, Ontario.
- Federici, A., & McMain, S. (2009). Recent advances in psychosocial treatments for borderline personality disorder. *Medicine Reports*, 1, 51.
- Koons, C. R., Robins, C. J., Tweed, J. L., Lynch, T. R., Gonzalez, A. M., Morse, J. Q., Bishop, G. K., Butterfield, M. I., & Bastian, L. A. (2001). Efficacy of dialectical behaviour therapy in women veterans with borderline personality disorder. *Behaviour Therapy*, 32, 371-390.
- Linehan, M. M., Armstrong, H. E., Suarez, A., Allmon, D., & Heard, H. L. (1991). Cognitive-behavioural treatment of chronically parasuicidal borderline patients. *Archives of General Psychiatry*, 48, 1060-1064.
- Linehan, M. M., Comtois, K. A., Murray, A. M., Brown, M. Z., Gallop, R. J., Heard, H. L., Korsland, K. E., Tutek, D. A., Reynolds, S. K., & Lindenboim, N. (2006). Two-year randomized trial and follow-up of Dialectical Behaviour Therapy vs therapy by experts for suicidal behaviours and borderline personality disorder. *Archives of General Psychiatry*, 63, 757-766.
- Linehan, M. M., Dimeff, L. A., Reynolds, S. K., Comtois, K., Welch, S. S., Heagerty, P., & Kivlahan, D. R. (2002). Dialectical behaviour therapy versus comprehensive validation therapy plus 12-step for the treatment of opioid dependent women meeting criteria for borderline personality disorder. *Drug and Alcohol Dependence*, 67, 13-26.
- Linehan, M. M., Heard, H. L., & Armstrong, H. E. (1993). Naturalistic follow-up of a behavioural treatment for chronically parasuicidal borderline patients. *Archives of General Psychiatry*, 50, 971-974.
- Linehan, M. M., Schmidt, H., Dimeff, L. A., Craft, J. C., Kanter, J. & Comtois, K. A. (1999). Dialectical Behaviour Therapy for patients with borderline personality disorder and drug-dependence. *American Journal of Addictions*, 8, 279-292.
- Linehan, M. M., Tutek, D. A., Heard, H. L., Armstrong, H. E. (1994). Interpersonal outcome of cognitive behavioural treatment for chronically suicidal borderline patients. *American Journal of Psychiatry*, 151, 1771-1776.
- Verheul, R., van den Bosch, L. M. C., Koeter, M. W. J., De Ridder, M. A. J., Stijnen, T., & van den Brink, W. (2003). Dialectical Behaviour Therapy for women with borderline personality disorder: 12 month, randomized clinical trial in The Netherlands. *British Journal of Psychiatry*, 182, 135-140.
- van den Bosch, L. M., Verheul, R., Schippers, G. M., & van den Brink, W. (2002). Dialectical Behavioural Therapy of borderline patients with and without substance abuse problems: Implementation and long-term effects. *Addictive Behaviour*, 27, 911-923.
- McMain, S. F., Guimond, T., Streiner, D. L., Cardish, R. J., Links, P. S. (2012). Dialectical Behaviour Therapy compared with general psychiatric management for borderline personality disorder: Clinical outcomes and functioning over a 2-year follow-up. *American Journal of Psychiatry*, 169, 650-661.
- Rathus, J. H., & Miller, A. L. (2002). Dialectical Behaviour Therapy adapted for suicidal adolescents. *Suicide and Life-Threatening Behaviour*, 32, 146-157.
- Harley, R., Sprich, S., Safran, S., Jacobo, M., & Fava, M. (2008). Adaptation of Dialectical Behaviour Therapy skills training group for treatment-resistant depression. *Journal of Nervous and Mental Disease*, 196(2), 136-143.
- Chen, E. Y., Matthews, L., Allan, C., Kuo, J. R., & Linehan, M. M. (2008). Dialectical Behaviour Therapy for clients with binge-eating disorder or bulimia nervosa and borderline personality disorder. *International Journal of Eating Disorders*, 41, 505-512.
- Axelrod, S. R., Perepletchikova, F., Holtzman, K., & Sinha, R. (2011). Emotion regulation and substance use frequency in women with substance dependence and borderline personality disorder receiving Dialectical Behaviour Therapy. *American Journal of Drug and Alcohol Abuse*, 37(1), 37-42.
- Harned, M. S., Korslund, K. E., Foa, E. B., & Linehan, M. M. (2012). Treating PTSD in suicidal and self-injuring women with borderline personality disorder: Development and preliminary evaluation of a Dialectical Behaviour Therapy Prolonged Exposure Protocol. *Behaviour Research and Therapy*, 50, 381-386.
- Harned, M. S., Jackson, S. C., Comtois, K. A., & Linehan, M. M. (2012). Dialectical Behaviour Therapy as a precursor to PTSD treatment for suicidal and self-injuring women with borderline personality disorder. *Journal of Traumatic Stress*, 23(4), 421-429.
- Neacsu, A. D., Rizvi, S. L., & Linehan, M. M. (2010). Dialectical Behaviour Therapy skills use as a mediator and outcome of treatment for borderline personality disorder. *Behaviour Research and Therapy*, 48, 832-839.
- Harley, R. M., Baity, M. R., Blais, M. A., & Jacobo, M. C. (2007). Use of Dialectical Behaviour Therapy skills training for borderline personality disorder in a naturalistic setting. *Psychotherapy Research*, 17, 362-370.
- Sambrook, S., Abba, N., Chadwick, P. (2006). Evaluation of a DBT emotional coping skills group for people with parasuicidal behaviours. *Behavioural and Cognitive Psychotherapy*, 35, 241-244.
- Safer, D. L., Telch, C. F., & Agras, W. S. (2001). Dialectical Behaviour Therapy for bulimia nervosa. *American Journal of Psychiatry*, 158, 632-634.
- Gratz, K. L., & Gunderson, J. G. (2006). Preliminary data on an acceptance-based emotion regulation group intervention for deliberate self-harm among women with borderline personality disorder. *Behaviour Therapy*, 37, 25-35.

A Positive Psychology Approach to Education for At-Risk High School Youth

Sophia Fanourgiakis, Ph.D. student and Margaret Lumley, Ph.D., C.Psych., University of Guelph

References

- Franklin, C., Streeter, C. L., Kim, J. S., & Tripodi, S. J. (2007). The effectiveness of a solution-focused, public alternative school for dropout prevention and retrieval. *Children & Schools*, 29(3), 133-144.
- Rumberger, R. W. (1987). High school dropouts: A review of issues and evidence. *Review of Educational Research*, 57(2), 101-121.
- Lerner, R. M., Lerner, J. V., Almerigi, J., & Theokas, C. (2005). Positive youth development: A view of the issues. *Journal of*



- Early Adolescence*, 25(1), 10-16.
4. Rawana, E., & Brownlee, K. (2009). Making the possible probable: A strength-based assessment and intervention framework for clinical work with parents, children and adolescents. *Families in Society*, 90(3), 255-260.
 5. Baker, J. A. (1998). The social context of school satisfaction among urban, low-income, AfricanAmerican students. *School Psychology Quarterly*, 13(1), 25-44.
 6. Murray, C. & Murray, K. M. (2004). Child level correlates of teacher-student relationships: An examination of demographic characteristics, academic orientations, and behavioral orientations. *Psychology in the Schools*, 41(7), 751-762.

Treating Insomnia in Primary Care: A Key Role for Psychologists

Judith R. Davidson, Ph.D., C. Psych., Kingston Family Health Team and Departments of Psychology and Oncology, Queen's University, Kingston, Ontario, Canada

References

1. Aikens JE, Rouse ME. Help-seeking for insomnia among adult patients in primary care. *Journal of the American Board of Family Practice* 2005;18: 257-261.
2. Simon GE, VonKorff M. Prevalence, burden, and treatment of insomnia in primary care. *American Journal of Psychiatry* 1997;154(10): 1417-1423.
3. Terzano MG, Parrino L, Cirignotta F, Ferini-Strambi L, Gigli G, Rudelli G. Studio Mofeo: insomnia in primary care, a survey conducted on the Italian population. *Sleep Medicine* 2004; 5: 67-75.
4. Davidson JR, Aimé A, Ivers H, Morin CM. Characteristics of individuals with insomnia who seek treatment in a clinical setting versus those who volunteer for a randomized controlled trial. *Behavioral Sleep Medicine* 2009;7: 37-52.
5. Baglioni C, Battagliese G, Feige B et al. Insomnia as a predictor of depression: a meta-analytic evaluation of longitudinal epidemiological studies. *Journal of Affective Disorders* 2011; 135(1-3): 10-19.
6. Daley M, Morin CM, LeBlanc M, Gregoire JP, Savard J, Bailleux L. Insomnia and its relationship to health-care utilization, work absenteeism, productivity and accidents. *Sleep Medicine* 2009;10(4): 427-438.
7. Hayward R, Jordan KP, Croft P. Healthcare use in adults with insomnia: a longitudinal study. *British Journal of General Practice* 2010; 60: 334-340.
8. Siriwardena AN, Apekey T, Tilling M, Dyas, JV, Middleton H, Orner R. General practitioners' preferences for managing insomnia and opportunities for reducing hypnotic prescribing. *Journal of Evaluation in Clinical Practice* 2010; 16: 731-737.
9. Morgenthaler R, Kramer M, Alessi C, Friedman L, Boehlecke B, Brown T, et al. Practice parameters for the psychological and behavioural treatment of insomnia: An update. An American Academy of Sleep Medicine Report. *Sleep* 2006;29(11): 1415-1419.
10. Schutte-Rodin S, Broch L, Buysse D, Dorsey C, Sateia M. Clinical guideline for the evaluation and management of chronic insomnia in adults. *Journal of Clinical Sleep Medicine* 2008;4(5): 487-504.
11. Wilson SJ, Nutt DJ, Alford C, Argyropoulos SV, Baldwin DS, Bateson AN et al. British Association for Psychopharmacology consensus statement on evidence-based treatment of insomnia, parasomnias and circadian rhythm disorders. *J Psychopharmacol*. 2010; 24(11): 1577-1600.
12. Smith MT, Huang MI, Manber R. Cognitive behavior therapy for chronic insomnia occurring within the context of medical and

- psychiatric disorders. *Clinical Psychology Review* 2005;25(5): 559-592.
13. Manber R, Edinger JD, Gress JL, San Pedro-Salcedo MG, Kuo TF, Kalista T. Cognitive behavioral therapy for insomnia enhances depression outcome in patients with comorbid major depressive disorder and insomnia. *Sleep* 2008;31(4): 489-495.
 14. Vitiello MV, Rybarczyk B, VonKorff M, Stepanski EJ. Cognitive behavioral therapy for insomnia improves sleep and decreases pain in older adults with co-morbid insomnia and osteoarthritis. *Journal of Clinical Sleep Medicine* 2009;5: 355-362.
 15. Morin CM, Colecchi C, Stone J, Sood R, Brink D. Behavioral and pharmacological therapies for late-life insomnia. A randomized controlled trial. *JAMA* 1999; 281(11): 991-999.
 16. Espie CA, Inglis SJ, Tessier S, Harvey L. The clinical effectiveness of cognitive behaviour therapy for chronic insomnia: implementation and evaluation of a sleep clinic in general medical practice. *Behaviour Research and Therapy* 2001;39: 45-60.
 17. Bastien CH, Vallieres A, Morin CM. Validation of the Insomnia Severity Index as an outcome measure for insomnia research. *Sleep Medicine* 2001;2(4): 297-307.
 18. Bootzin, R. R. Stimulus control treatment for insomnia. *Proceedings of the American Psychological Association* 1972; 7: 395-396.
 19. Spielman, A. J., Saskin, P., & Thorpy, M. J. Treatment of chronic insomnia by restriction of time in bed. *Sleep* 1987; 10: 45-56.
 20. Davidson JR. Treating insomnia in primary care -Early recognition and management. *Insomnia Rounds* 2012 1(3): 1-10. www.insomniarounds.ca

Providing Psychological Services and Training in Primary Care: Ryerson University and St. Michael's Hospital

Jen Rouse, MA, Kelly McShane, PhD, & Candice Monson, PhD Ryerson University

References

1. Smetanin, P., Stiff, D., Briante, C., Adair, C., Ahmad, S. & Khan, M. (2011). The life and economic impact of major mental illnesses in Canada: 2011 to 2041. Risk Analytica, on behalf of the Mental Health Commission of Canada.
2. Chambliss, D. L., & Ollendick, T. H. (2001). Empirically supported psychological interventions: Controversies and evidence. *Annual Review of Psychology*, 52, 685-716. doi: 10.1146/annurev.psych.52.1.685
3. Hunsley, J. (2002). *The cost-effectiveness of psychological interventions*. Retrieved from <http://www.cpa.ca/cpasite/userfiles/Documents/publications/Cost-Effectiveness.pdf>
4. Myhr, G., & Payne, K. (2006). Cost-effectiveness of cognitive-behavioural therapy for mental disorders: Implication for public health care funding policy in Canada. *Canadian Journal of Psychiatry*, 51, 662-670.
5. United Kingdom Department of Health. (2001). *Treatment choice in psychological therapies and counselling: Evidence based clinical practice guidelines*. Retrieved from http://www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/@dh/@en/documents/digitalasset/dh_4058245.pdf
6. Mental Health Commission of Canada. (2012a). *Changing directions, changing lives: The mental health strategy for Canada*. Calgary, AB: Author. Retrieved from: strategy.mentalhealthcommission.ca/pdf/strategy-images-en.pdf
7. Health Care Innovation Working Group. (2012). *From innovation to action: The first report of the health care working group*. Retrieved from: www.councilofthefederation.ca/pdfs/Health%20Innovation%20Report-E-WEB.pdf



**Transdiagnostic Group Cognitive-Behavioral Therapy:
An Efficacious and Cost-Effective Method Across Diverse
Populations**

Authors: Backs-Dermott, Barbara Ph.D.,^{1,2}; Hamilton, Kate Ph.D.,¹; Macrodimitris, Sophie, Ph.D.,^{1,2}; Mothersill, Kerry, Ph.D.,^{1,2}; Garinger, Jennifer Ph.D.,¹; Luu, May B.A. Honours,²; Wershler, Julie B.A. Honours,³; & Ching, Laurie M.Sc.,^{1,2}, 1. Alberta Health Services, 2. University of Calgary, 3. University of New Brunswick

References

1. Canadian Psychiatric Association (2006). Clinical practice guidelines: Management of anxiety disorders. *The Canadian Journal of Psychiatry / La Revue canadienne de psychiatrie*, 51(Supplement 2), 1S-92S.
2. National Institute for Clinical Excellence (2007). *Depression (amended): Management of depression in primary and secondary care*. London, UK: National Institute for Health and Clinical Excellence.
3. Stewart, R. E., & Chambless, D. L. (2009). Cognitive-behavioral therapy for adult anxiety disorders in clinical practice: A meta-analysis of effectiveness studies. *Journal of Consulting and Clinical Psychology*, 77(4), 595-606.
4. Royal Australian and New Zealand College of Psychiatrists Clinical Practice Guidelines Team for Panic Disorder and Agoraphobia, T. (2003). Australian and New Zealand clinical practice guidelines for the treatment of panic disorder and agoraphobia. *Australian and New Zealand Journal of Psychiatry*, 37(6), 641-656.
5. Vos, T., Corry, J., Haby, M. M., Carter, R., & Andrews, G. (2005). Cost-effectiveness of cognitive-behavioural therapy and drug interventions for major depression. *Australian and New Zealand Journal of Psychiatry*, 39(8), 683-692.
6. Shafran, R., Clark, D., Fairburn, C., Arntz, A., Barlow, D., Ehlers, A., et al. (2009). Mind the gap: Improving the dissemination of CBT. *Behaviour Research and Therapy*, 47(11), 902-909.
7. Barlow, D. H. (Ed.). (2001). *Clinical handbook of psychological disorders. A step-by-step treatment manual* (3rd ed.). New York: Guilford.
8. Bourne, E. J. (1998). *Overcoming specific phobia: A hierarchy and exposure-based protocol for the treatment of all specific phobias*. Oakland, CA: New Harbinger.
9. Barlow, D. H., & Craske, M.G. (2000). *Mastery of your anxiety and panic* (third edition). New York: Oxford University Press.
10. Emery, G. (2000). *Overcoming depression: A cognitive-behavior protocol for the treatment of depression*. Oakland, CA: New Harbinger.
11. Beck, A. T., Rush, A. J., Shaw, B. F., & Emery, G. (1979). *Cognitive therapy of depression*. New York: Guilford.
12. Beck, J. (1995). *Cognitive therapy: Basics and beyond*. New York: Guilford.
13. Bourne, E. J. (2000). *The anxiety and phobia workbook*. New York: New Harbinger.
14. Burns, D. D. (1980). *Feeling good*. New York: Signet.
15. Burns, D. D. (1989). *The Feeling Good Handbook*. New York: William Morrow and Co.
16. Dobson, K. S. (2001). *Handbook of cognitive-behavioral therapies* (2nd ed.). New York: Guilford.
17. Greenberger, D., & Padesky, C. A. (1995). *Mind over mood*. New York: Guilford.
18. Kabat-Zinn, J., & Santorelli, S. (1993). *Mindfulness-based stress reduction professional training resource manual*. Worcester, MA: University of Massachusetts Medical Center.
19. Macrodimitris, S.D., Hamilton, K.E., Backs-Dermott, B.J., & Mothersill, K.J. (2010). CBT Basics: A group approach to teaching fundamental cognitive-behavioral skills. *Journal of Cognitive Psychotherapy: An International Quarterly*, 24(2), 132-146.
20. Hamilton, K., & Macrodimitris, S. (2008). *CBT Basics II: Participant workbook*. Calgary, AB, Canada: Alberta Health Services.
21. Macrodimitris, S. D., & Hamilton, K. (2008). *CBT Basics II: Therapist manual*. Calgary, AB, Canada: Alberta Health Services.
22. Teasdale, J. D., Moore, R. G., Hayhurst, H., Pope, M., Williams, S., & Segal, Z. V. (2002). Metacognitive awareness and prevention of relapse in depression: Empirical evidence. *Journal of Consulting and Clinical Psychology*, 70(2), 275-287.
23. Hofmann, S.G., Sawyer, A.T., Witt, A.A., & Oh, D. (2010). The effects of mindfulness-based therapy on anxiety and depression: A meta-analytic review. *Journal of Consulting and Clinical Psychology*, 78(2), 169-183.
24. Hamilton, K.E., Wershler, J.L., Macrodimitris, S.D., Backs-Dermott, B.J., Ching, L.E., & Mothersill, K.J. (2012). Exploring the effectiveness of a mixed-diagnosis group cognitive-behavioral therapy intervention across diverse populations. *Cognitive and Behavioral Practice*, 19(3), 472-482.